

Invariant manifold of your dynamical system

A. J. Roberts, University of Adelaide
<http://orcid.org/0000-0001-8930-1552>

11:47am, November 25, 2022

Throughout and generally: the lowest order, most important, terms are near the end of each expression.

The specified dynamical system

$$\dot{u}_1 = u_3$$

$$\dot{u}_2 = u_4$$

$$\dot{u}_3 = f_1 \varepsilon^2 (1/2 \exp(-it)i - 1/2 \exp(it)i) + \varepsilon^2 (-1/2 u_1^3 - 3/10 u_3 + 3/10 u_4) - 2u_1 + u_2$$

$$\dot{u}_4 = f_2 \varepsilon^2 (3/20 \exp((-it)/2)u_3 - 3/10 \exp((-it)/2)u_4 + 3/20 \exp((it)/2)u_3 - 3/10 \exp((it)/2)u_4) + u_1 - 2u_2$$

Invariant subspace basis vectors

$$\vec{e}_1 = \{ \{1, 1, i, i\}, \exp(it) \}$$

$$\vec{e}_2 = \{ \{1, 1, -i, -i\}, \exp(-it) \}$$

$$\vec{z}_1 = \{ \{1/4, 1/4, 1/4i, 1/4i\}, \exp(it) \}$$

$$\vec{z}_2 = \{ \{1/4, 1/4, -1/4i, -1/4i\}, \exp(-it) \}$$

off echo;

The invariant manifold These give the location of the invariant manifold in terms of parameters s_j .

$$u_1 = f_1 \varepsilon^2 (3/16 \exp(-it)i - 3/16 \exp(it)i) + f_2 \varepsilon^2 (4/55 \exp(-it/2)s_2i - 4/25 \exp(-3it/2)s_2i - 4/55 \exp(it/2)s_1i + 4/25 \exp(3it/2)s_1i) + \varepsilon^2 (-9/16 \exp(-it)s_2^2s_1 + 7/96 \exp(-3it)s_2^3 - 9/16 \exp(it)s_2s_1^2 + 7/96 \exp(3it)s_1^3) + \exp(-it)s_2 + \exp(it)s_1 + O(\varepsilon^4)$$

$$u_2 = f_1 \varepsilon^2 (-1/16 \exp(-it)i + 1/16 \exp(it)i) + f_2 \varepsilon^2 (7/55 \exp(-it/2)s_2i + 1/25 \exp(-3it/2)s_2i - 7/55 \exp(it/2)s_1i - 1/25 \exp(3it/2)s_1i) + \varepsilon^2 (3/16 \exp(-it)s_2^2s_1 - 1/96 \exp(-3it)s_2^3 + 3/16 \exp(it)s_2s_1^2 - 1/96 \exp(3it)s_1^3) + \exp(-it)s_2 + \exp(it)s_1 + O(\varepsilon^4)$$

$$u_3 = f_1 \varepsilon^2 (1/16 \exp(-it) + 1/16 \exp(it)) + f_2 \varepsilon^2 (2/55 \exp(-it/2)s_2 - 6/25 \exp(-3it/2)s_2 + 2/55 \exp(it/2)s_1 - 6/25 \exp(3it/2)s_1) + \varepsilon^2 (3/16 \exp(-it)s_2^2s_1i - 7/32 \exp(-3it)s_2^3i - 3/16 \exp(it)s_2s_1^2i + 7/32 \exp(3it)s_1^3i) - \exp(-it)s_2i + \exp(it)s_1i + O(\varepsilon^4)$$

$$u_4 = f_1 \varepsilon^2 (-3/16 \exp(-it) - 3/16 \exp(it)) + f_2 \varepsilon^2 (7/110 \exp(-it/2)s_2 + 3/50 \exp(-3it/2)s_2 + 7/110 \exp(it/2)s_1 + 3/50 \exp(3it/2)s_1) + \varepsilon^2 (-9/16 \exp(-it)s_2^2s_1i + 1/32 \exp(-3it)s_2^3i + 9/16 \exp(it)s_2s_1^2i - 1/32 \exp(3it)s_1^3i) - \exp(-it)s_2i + \exp(it)s_1i + O(\varepsilon^4)$$

Invariant manifold ODEs The system evolves on the invariant manifold such that the parameters evolve according to these ODEs.

$$\dot{s}_1 = f_1 \varepsilon^4 (9/64 s_2 s_1 - 9/128 s_1^2 + 3/160i) - 1/8 f_1 \varepsilon^2 + 93/5500 f_2^2 \varepsilon^4 s_1i + \varepsilon^4 (-155/256 s_2^2 s_1^3i + 9/160 s_2 s_1^2) + 3/8 \varepsilon^2 s_2 s_1^2i + O(\varepsilon^5)$$

$$\dot{s}_2 = f_1 \varepsilon^4 (-9/128 s_2^2 + 9/64 s_2 s_1 - 3/160i) - 1/8 f_1 \varepsilon^2 - 93/5500 f_2^2 \varepsilon^4 s_2i + \varepsilon^4 (155/256 s_2^3 s_1^2i + 9/160 s_2^2 s_1) - 3/8 \varepsilon^2 s_2^2 s_1i + O(\varepsilon^5)$$